

Summary of feed diets influencing boar taint compounds

	Diet / Feed additive	Reduced Skatol content	Reduced Androstenon content
	Casein instead of yeast slurry	yes	
	Dietary Amino Acids	no	yes
	Reduce indigestible protein	no	no
	Lysine	no	
	Wet feed with whey vs. Dry feed	yes	
	Fructo-oligosaccharides (short chain)	yes	
	Crude Protein	yes	
	Finisher diet with 5% oligofructose	yes	
	Finisher diet with 5% inuline	yes	
chicory	Dried chichory root (10%)	no	yes
chicory	Dried chichory with 60% inulin (9%)	yes	
chicory	Chichory inuline (70% inuline)	yes	no
chicory	Chichory root (5%)	yes	no
chicory	Chichory root (15%)	yes	no
lupin	Lupin	no	yes
lupin	Lupin	yes	
patato	Pelleted diet with patato starch (20%)	no	no
patato	Raw patato starch (20%) as top dressing	yes	
patato	Patato starch supplement	yes	no
sugar beet	Sugar beet pulp	yes	no
sugar beet	Sugar beet pulp added tot yeast slurry feed	yes	
other	Palm cakes	yes	no
other	Jerusalem artichoke	yes	no
other	Taintstop	yes	yes
other	Pure grain	yes	no
other	Peas or faba beans	no	

Source: <http://www.ca-ipema.eu/> and <https://zenodo.org/record/1460483#.XY3THGAzaMo>